arXiv:0904.0570v2 [cs.LO] 26 Jun 2009

The Derivational Complexity Induced by the Dependency Pair Method*

Georg Moser and Andreas Schnabl
{georg.moser,andreas.schnabl}@uibk.ac.at

June 26, 2009

We investigate the derivational complexity induced by the (basic) dependency pair method based on reasonably strong base orders. Suppose the derivational complexity induced by a (direct) termination method is closed under elementary functions. Then we show that the derivational complexity induced by the dependency pair method based on this termination technique is the same as for the direct technique. In particular this implies that the derivational complexity induced by the dependency pair method based on lexicographic path orders or multiset path orders is multiple recursive or primitive recursive, respectively. Moreover for the dependency pair method based on Knuth-Bendix orders, we obtain that the derivational complexity function is majorised by the Ackermann function. These characterisations are essentially optimal.

---

* This research is partly supported by FWF (Austrian Science Fund) project P20133.

Contents

1 Introduction **2**

2 Dependency Pairs **4**

3 Progenitor and Progeny **5**

4 Dependency Pairs and Complexity **10**

5 The Lower Bound **15**

6 Dependency Graphs **16**

7 Conclusion **18**

1 Introduction

In order to assess the complexity of a (terminating) term rewrite system (TRS for short) it is natural to look at the maximal length of derivation sequences, as suggested by Hofbauer and Lautemann in [12]. More precisely, the *derivational complexity function* with respect to a (finite and terminating) TRS $\mathcal{R}$ relates the length of the longest derivation sequence to the size of the initial term. For direct termination methods a considerable number of results establishes (essentially optimal) upper bounds on the growth rate of the derivational complexity function. See e.g. [13, 16] for recent results in this direction. However, for transformation techniques like semantic labelling [21] or the dependency pair method [1] the situation changes drastically. It is a trivial observation that labelling with finite models exactly preserves the derivational complexity of a TRS. This obviously holds for infinite models, as well. However, it is not as easily applicable in this setting since the labelled TRS is infinite, so bounding the derivational complexity might be even harder for the labelled TRS than for the original TRS. Apart from this trivial case, only partial results are known. With respect to semantic labelling, [15] establishes bounds on the derivation length of TRS, when natural numbers are used as labels and termination is shown via the Knuth-Bendix order (KBO). And recently in [8, 9] the derivation length induced by the basic dependency pair method is investigated. Still in both cases only restricted variants of semantic labelling or the dependency pair method could be analysed, compare [15, 8, 9].

In this paper we investigate the derivational complexity induced by the (basic) dependency pair method based on reasonably strong base orders. Suppose the class of derivational complexity functions induced by a (direct) termination method is closed under elementary functions. Then we show that the derivational complexity induced by the dependency pair method based on this termination technique is the same as for the direct technique.

More precisely we show that the derivational complexity of a TRS whose termination is established via the dependency pair method combined with some base order is triple exponential in the derivational complexity induced by the base order directly. Moreover, we present an example which shows that at least two of the three exponentials in our upper bound can actually be reached.

It should be emphasised that the notion of dependency pair method studied here amounts to the original technique as introduced by Arts and Giesl [1] (see also [6]). Consider the following TRS $\mathcal{R}_1$ taken from [3] that we use in the sequel to illustrate the developed notions.

$$\begin{array}{llll} 1\colon & (x \times y) \times z \to x \times (y \times z) & 3\colon & (x + y) \times z \to (x \times z) + (y \times z) \\ 2\colon & z \times (x + \mathsf{f}(y)) \to \mathsf{g}(z, y) \times (x + \mathsf{a}) & & \end{array}$$

Due to rule 2, termination of $\mathcal{R}_1$ cannot be concluded by the lexicographic path order (LPO), cf. [3]. On the other hand, termination follows easily by the dependency pair method based on LPO, if we make use of argument filterings.

The gist of our result is that for this (standard) application of the dependency pair method the derivational complexity induced by LPO bounds the derivation lengths admitted by the investigated TRS $\mathcal{R}_1$. From this we can conclude that the derivational complexity function with respect to $\mathcal{R}_1$ is at most a multiple recursive function. (Recall that the derivational complexity induced by LPO is multiple recursive, cf. [20].) An analogous result holds if we employ the multiset path order (MPO) or KBO as base order. Moreover the thus obtained upper bound is still tight, which essentially follows from the tight characterisation of the derivational complexity of the indicated base orders [10, 20, 14].

Notice the challenges of such an investigation: In order to estimate the derivation length of $\mathcal{R}_1$ we only consider the derivation length induced by the base order. This implies that we use an upper bound on the maximal number of dependency pair steps to bound the length of derivations. It remains open to what extent such a result holds in general. The challenge of such an endeavour is most prominent if we allow an iterative use of the dependency pair transformation as for example in the recursive SCC algorithm (see [6]) or in the dependency pair framework (see [19]). It is well-known that two iterations of the recursive SCC algorithm based on the subterm criterion (see [7]) suffice to show termination of (the standard formulation) of the Ackermann function. See also [4] for a like minded example. Clearly in this context a triple exponential function is by far not sufficient to bound the difference between the derivational complexity of the TRS and the derivational complexity induced by the base method.

The rest of this paper is organised as follows. In Section 2 we present basic notions and starting points of the paper. Section 3 introduces suitable notions to trace an (implicit) dependency pair derivation in a given derivation over some TRS $\mathcal{R}$. Our main result is proved in Section 4, while Section 5 presents the above mentioned example on the lower bound of the correspondence. Finally, we conclude in Section 7.

2 Dependency Pairs

We assume familiarity with the basics of term rewriting, see [2, 18]. Below we recall the bare essentials of the (basic) dependency pair method as put forward in [1], but (at least nodding) acquaintance with [1] or [6] will prove helpful.

Let $\mathcal{V}$ denote a countably infinite set of variables and $\mathcal{F}$ a signature. The set of terms over $\mathcal{F}$ and $\mathcal{V}$ is denoted by $\mathcal{T}(\mathcal{F},\mathcal{V})$. The (proper) subterm relation is denoted as $\trianglelefteq$ ($\triangleleft$). The *root symbol* (denoted as $\mathsf{rt}(t)$) of a term t is either t itself, if $t \in \mathcal{V}$, or the symbol f, if $t = f(t_1,\ldots,t_n)$. The *set of positions* $\mathcal{P}\mathsf{os}(t)$ of a term t is defined as usual. We write $p \leqslant q$ ($p < q$) to denote that p is a (proper) prefix of q, and $p \parallel q$ if neither $p \leqslant q$ nor $q \leqslant p$. The subterm of t at position p is denoted as $t|_p$. We write $\mathcal{P}\mathsf{os}_{\mathcal{F}}(t)$ ($\mathcal{P}\mathsf{os}_{\mathcal{V}}(t)$) for the set of positions p such that $\mathcal{F}$ ($\mathcal{V}$) contains $\mathsf{rt}(t_p)$. The *size* $|t|$ and the *depth* $\mathsf{dp}(t)$ of a term t are defined as usual (e.g., $|\mathsf{f}(\mathsf{a},x)| = 3$ and $\mathsf{dp}(\mathsf{f}(\mathsf{a},x)) = 1$). To simplify the exposition, we often confuse terms and their tree representations. I.e., we call a maximal set of positions B in a term t such that for no $q, q' \in B$, we have $q \parallel q'$, a *branch* of t.

Let $\mathcal{R}$ be a finite TRS over $\mathcal{F}$. We write $\to_{\mathcal{R}}$ (or simply $\to$) for the induced rewrite relation. If we wish to indicate the redex position p and the applied rewrite rule $l \to r$ in a reduction from s to t, we write $s \to_{p,l\to r} t$. The set of defined function symbols is denoted as $\mathcal{D}$, while the constructor symbols are collected in $\mathcal{C}$. The n-fold composition of $\to$ is denoted as $\to^n$ and the *derivation length* of a terminating term s with respect to a rewrite relation $\to$ is defined as $\mathsf{dl}(s,\to) := \max\{n \mid \exists t\ s \to^n t\}$. The *derivational complexity function* (with respect to a TRS $\mathcal{R}$) is defined as: $\mathsf{dc}_{\mathcal{R}}(n) = \max\{\mathsf{dl}(t,\to_{\mathcal{R}}) \mid |t| \leqslant n\}$.

In analogy to the derivational complexity function $\mathsf{dc}_{\mathcal{R}}$ we define functions tracing the size or depth. The *potential size* of a terminating term s with respect to a relation $\to$ is defined as follows: $\mathsf{psz}(s,\to) := \max\{|t| \mid s \to^* t\}$ and the induced *size growth function* (with respect to $\mathcal{R}$) is defined as $\mathsf{szg}_{\mathcal{R}}(n) := \max\{\mathsf{psz}(t,\to_{\mathcal{R}}) \mid |t| \leqslant n\}$. The *potential depth* $\mathsf{pdp}(s,\to)$ of a terminating term s and the *depth growth function* $\mathsf{dpg}_{\mathcal{R}}(n)$ are defined similarly.

We recall the central notions of the dependency pair method, see [1, 6]. Let t be a term. We set $t^\sharp := t$ if $t \in \mathcal{V}$, and $t^\sharp := f^\sharp(t_1,\ldots,t_n)$ if $t = f(t_1,\ldots,t_n)$. Here $f^\sharp$ is a new n-ary function symbol called *dependency pair symbol*. For a signature $\mathcal{F}$, we define $\mathcal{F}^\sharp = \mathcal{F} \cup \{f^\sharp \mid f \in \mathcal{F}\}$. The set $\mathsf{DP}(\mathcal{R})$ of *dependency pairs* of a TRS $\mathcal{R}$ is defined as $\{l^\sharp \to u^\sharp \mid l \to r \in \mathcal{R}, u \trianglelefteq r, \mathsf{rt}(u) \in \mathcal{D}, l \not\triangleright u\}$.

Proposition 2.1. *A TRS $\mathcal{R}$ is terminating if and only if there exists no infinite derivation of the form $t_1^\sharp \to_{\mathcal{R}}^* t_2^\sharp \to_{\mathsf{DP}(\mathcal{R})} t_3^\sharp \to_{\mathcal{R}}^* \ldots$ such that for all $i > 0$, $t_i^\sharp$ is terminating with respect to $\mathcal{R}$.*

Proposition 2.1 gives rise to the *dependency pair complexity function*

$$\mathsf{DPc}_{\mathcal{R}}(n) := \max\{\mathsf{dl}(t^\sharp, \to_{\mathsf{DP}(\mathcal{R})/\mathcal{R}}) \mid |t| \leqslant n\}\ ,$$

where we write $\to_{\mathsf{DP}(\mathcal{R})/\mathcal{R}}$ for $\to_{\mathcal{R}}^* \cdot \to_{\mathsf{DP}(\mathcal{R})} \cdot \to_{\mathcal{R}}^*$, cf. [5]. Now, we fix the notion of *basic* dependency pair method. An *argument filtering* (for a signature $\mathcal{F}$) is a mapping π that assigns to every n-ary function symbol $f \in \mathcal{F}$ an argument position $i \in \{1,\ldots,n\}$ or a (possibly empty) list $[i_1, \dots, i_m]$ of argument positions with $1 \leqslant i_1 < \dots < i_m \leqslant n$. The signature $\mathcal{F}_\pi$ consists of all function symbols f such that $\pi(f)$ is some list $[i_1, \dots, i_m]$, where in $\mathcal{F}_\pi$ the arity of f is m. Every argument filtering π induces a mapping from $\mathcal{T}(\mathcal{F}, \mathcal{V})$ to $\mathcal{T}(\mathcal{F}_\pi, \mathcal{V})$, also denoted by π:

$$\pi(t) = \begin{cases} t & \text{if } t \text{ is a variable} \\ \pi(t_i) & \text{if } t = f(t_1, \dots, t_n) \text{ and } \pi(f) = i \\ f(\pi(t_{i_1}), \dots, \pi(t_{i_m})) & \text{if } t = f(t_1, \dots, t_n) \text{ and } \pi(f) = [i_1, \dots, i_m] \end{cases}$$

A *reduction pair* $(\gtrsim, \succ)$ consists of a rewrite preorder $\gtrsim$ and a compatible well-founded order $\succ$ which is closed under substitutions. Here compatibility means the inclusion $\gtrsim \cdot \succ \cdot \gtrsim \subseteq \succ$.

Proposition 2.2. *A TRS $\mathcal{R}$ is terminating if and only if there exist an argument filtering π and a reduction pair $(\gtrsim, \succ)$ such that $\pi(\mathsf{DP}(\mathcal{R})) \subseteq \succ$ and $\pi(\mathcal{R}) \subseteq \gtrsim$.*

Let $\mathcal{R}$ be a terminating TRS. In the sequel we show that the derivational complexity function $\mathsf{dc}_\mathcal{R}$ is bounded triple exponentially in the dependency pair complexity function $\mathsf{DPc}_\mathcal{R}$. For that we mainly bound the depth growth function of $\mathcal{R}$ exponentially in $\mathsf{DPc}_\mathcal{R}$. As the maximal length of a derivation is exponentially bounded in the size of the occurring terms and the latter is exponentially bounded in their depth, our result then follows. In order to prove the main step we analyse the shape of a potential derivation over $\mathcal{R} \cup \mathsf{DP}(\mathcal{R})$ in the light of a given $\mathcal{R}$-derivation. This is the purpose of the next section.

3 Progenitor and Progeny

To analyse the shape of a potential derivation over $\mathcal{R} \cup \mathsf{DP}(\mathcal{R})$ we introduce a specific generalisation of the notion of *descendant* of a position p which we call *progeny*. Recall the definition of descendants (see [18, Chapter 4]). Let $A : s \to_{p', l \to r} t$ be a rewriting step, and let $p \in \mathcal{P}\mathsf{os}(s)$. Then the *descendants of p in t* (denoted by $p \setminus A$) are defined as follows:

$$p \setminus A = \begin{cases} \{p\} & \text{if } p < p' \text{ or } p \parallel p', \\ \{p' q_3 q_2 \mid r|_{q_3} = l|_{q_1}\} & \text{if } p = p' q_1 q_2 \text{ with } q_1 \in \mathcal{P}\mathsf{os}_\mathcal{V}(l), \\ \varnothing & \text{otherwise} \end{cases}$$

In our context, we also want to keep track of redex positions, not just of positions in the context or the substitution of the rewrite rule. This intuition is cast into the following definition.

Definition 3.1. Let $A : s \to_{p', l \to r} t$ be a rewriting step, and let $p \in \mathcal{P}\mathsf{os}(s)$. Then the



Figure 1: A derivation, its progeny relation and redex positions.

progenies of p *in* t (denoted by $p \setminus\!\!\setminus A$) are:

$$p \setminus\!\!\setminus A = \begin{cases} \{p\} & \text{if } p < p' \text{ or } p \parallel p', \\ \{p'q_3q_2 \mid r|_{q_3} = l|_{q_1}\} & \text{if } p = p'q_1q_2 \text{ with } q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{V}}(l), \\ \{p'q_2 \mid r|_{q_2} = l|_{q_1}\} & \text{if } p = p'q_1 \text{ with } q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(l) \setminus \{\epsilon\}, \\ \{pq_1 \mid r|_{q_1} \not\trianglelefteq l \wedge q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)\} & \text{if } p = p' \end{cases}$$

If $q \in p \setminus\!\!\setminus A$, then we also say that p is a *progenitor of* q *in* s. We denote the set of progenitors of q in s by $A \setminus\!\!\setminus q$. For a set $P \subseteq \mathcal{P}\mathsf{os}(s)$, we define $P \setminus\!\!\setminus A = \bigcup_{p \in P} p \setminus\!\!\setminus A$.

Notice that the distinction between the last two cases corresponds to the exclusion of rules $l^\sharp \to u^\sharp$ from $\mathsf{DP}(\mathcal{R})$ where $u \triangleleft l$, see Section 2.

Example 3.1. Consider the TRS $\mathcal{R}_1$ from Section 1 and let $t_1 = (x \times y) \times (z + \mathsf{f}(w))$, $t_2 = x \times (y \times (z + \mathsf{f}(w)))$, and $t_3 = x \times (\mathsf{g}(y, w) \times (z + \mathsf{a}))$. We have the derivation $A : t_1 \to t_2 \to t_3$, cf. Figure 1. Redex positions are marked by circles, the progeny relation is marked by dotted and dashed lines (the lines will be distinguished in Example 3.4 below). For clarity progeny relations between variables have been omitted.

Lemma 3.1. *Let* $\mathcal{R}$ *be a TRS, let* $A : s \to_{\mathcal{R}} t$*, let* $p \in \mathcal{P}\mathsf{os}(s)$*, and let* $q \in \mathcal{P}\mathsf{os}(t)$*. If* $q \in p \setminus\!\!\setminus A$ *and* $\mathsf{rt}(t|_q) \in \mathcal{D}$*, then* $\mathsf{rt}(s|_p) \in \mathcal{D}$ *and* $(s|_p)^\sharp \to^{=}_{\mathcal{R} \cup \mathsf{DP}(\mathcal{R})} (t|_q)^\sharp$*.*

Proof. Suppose that A is $s \to_{p', l \to r} t$. If $p < p'$ or $p \parallel p'$, then by definition, we have $p = q$ and thus $(s|_p)^\sharp \to^{=}_{\mathcal{R}} (t|_q)^\sharp$. On the other hand, if $p = p'$, then there exists $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$ such that $q = p'q_1$. Moreover, $t|_q \not\trianglelefteq s|_p$. By assumption $\mathsf{rt}(t|_q) \in \mathcal{D}$ and thus we obtain $(s|_p)^\sharp \to_{\mathsf{DP}(\mathcal{R})} (t|_q)^\sharp$. Finally, if $p > p'$, then by definition, we have $s|_p = t|_q$. Then (trivially) $(s|_p)^\sharp \to^{=}_{\mathcal{R}} (t|_q)^\sharp$. $\square$

Lemma 3.2. *Let* $A : s \to_{p',l\to r} t$ *be a rewriting step. Then for every* $q \in \mathcal{P}\mathsf{os}(t)$*, we have* $A \setminus\!\!\setminus q \neq \emptyset$*.*

Proof. If $q < p'$ or $q \parallel p'$, then $A \setminus\!\!\setminus q = \{q\}$. If $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $t|_q \ntriangleleft s|_{p'}$, then $A \setminus\!\!\setminus q = \{p'\}$. If $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $t|_q \triangleleft s|_{p'}$, then there is some p_1 such that $s|_{p'p_1} = t|_q$, so $p'p_1 \in A \setminus\!\!\setminus q$. Last, if $q = p'q_1q_2$ and $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{V}}(r)$, then there is some p_1 such that $s|_{p'p_1} = t|_{p'q_1}$. Therefore, $p'p_1q_2 \in A \setminus\!\!\setminus q$. ☐

Definition 3.1 and Lemmata 3.1 and 3.2 extend to derivations in the natural way:

Definition 3.2. Let $A : s \to^* t$ be a derivation, and let $p \in \mathcal{P}\mathsf{os}(s)$. Then the *progenies of* p *in* t (also denoted by $p \setminus\!\!\setminus A$) are defined as follows:

- If A is the empty derivation, then $p \setminus\!\!\setminus A = \{p\}$.
- Otherwise, we can split A into $A_1 : s \to s'$ and $A_2 : s' \to^* t$. Then $p \setminus\!\!\setminus A = (p \setminus\!\!\setminus A_1) \setminus\!\!\setminus A_2$.

We say p is a *progenitor* of q if $p \in A \setminus\!\!\setminus q$, which holds if $q \in p \setminus\!\!\setminus A$. Moreover, we have $q \in P \setminus\!\!\setminus A$ if and only if $q \in p \setminus\!\!\setminus A$ for some $p \in P$.

The next lemma follows by straightforward induction using Lemmata 3.2 and 3.1.

Lemma 3.3. *Let* $A : s \to^* t$ *be a derivation, and let* $p \in \mathcal{P}\mathsf{os}(s)$*,* $q \in \mathcal{P}\mathsf{os}(t)$*. Then the set* $A \setminus\!\!\setminus q$ *of progenitors of* q *is not empty. Moreover if* $q \in p \setminus\!\!\setminus A$ *with* $\mathsf{rt}(t|_q) \in \mathcal{D}$*, then* $\mathsf{rt}(s|_p) \in \mathcal{D}$ *and* $(s|_p)^\sharp \to^*_{\mathcal{R}\cup\mathsf{DP}(\mathcal{R})} (t|_q)^\sharp$*.*

Using Lemma 3.3, we can extract derivations over $\mathcal{R} \cup \mathsf{DP}(\mathcal{R})$ from a given derivation in a TRS $\mathcal{R}$ using positions connected by the progeny relation.

Definition 3.3. Let $\mathcal{R}$ be a TRS, let $t_1, \ldots, t_n$ be terms, and let $p_1, \ldots, p_n$ be positions in $t_1, \ldots, t_n$, respectively, such that $\mathsf{rt}(t_n|_{p_n}) \in \mathcal{D}$, and for all $1 \leqslant i \leqslant n-1$, we have $A_i : t_i \to_{\mathcal{R}} t_{i+1}$ and $p_{i+1} \in p_i \setminus\!\!\setminus A_i$. Then we call $A : (t_1|_{p_1})^\sharp \to^*_{\mathcal{R}\cup\mathsf{DP}(\mathcal{R})} (t_n|_{p_n})^\sharp$ the *implicit dependency pair derivation* with respect to $t_1, \ldots, t_n$ and $p_1, \ldots, p_n$. We denote the number of $\mathsf{DP}(\mathcal{R})$-steps in A as $\mathsf{DPI}(A)$.

Notice that by Lemma 3.3, any implicit dependency pair derivation based on a derivation over $\mathcal{R}$ is indeed a derivation over $\mathcal{R} \cup \mathsf{DP}(\mathcal{R})$.

Example 3.2 (continued from Example 3.1)**.** The implicit dependency pair derivation with respect to t_1, t_2, t_3 and $\epsilon, 2, 2$ is given as follows:

$$t_1^\sharp \to_{\mathsf{DP}(\mathcal{R})} y \times^\sharp (z + \mathsf{f}(w)) \to_{\mathsf{DP}(\mathcal{R})} \mathsf{g}(y, w) \times^\sharp (z + \mathsf{a}) \,.$$

Lemma 3.4. *Let* $A : s \to_{p',l\to r} t$ *be a rewriting step. Let* $q, q' \in \mathcal{P}\mathsf{os}(t)$*. If* $q \leqslant q'$*, then for any* $p_0 \in A \setminus\!\!\setminus q$*, there exists* $p_0' \in A \setminus\!\!\setminus q'$ *such that* $p_0 \leqslant p_0'$*.*

Proof. According to Definition 3.1, there are four cases for q'.

- If $q' < p'$ or $q' \parallel p'$, then also $q < p'$ or $q \parallel p'$. Therefore, $A \mathbin{\backslash\!\!\backslash} q = \{q\}$ and $A \mathbin{\backslash\!\!\backslash} q' = \{q'\}$.
- If $q' = p'q'_1$, $q'_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q'_1} \ntriangleleft l$, then either $q < p'$, or $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q_1} \ntriangleleft l$. We have $A \mathbin{\backslash\!\!\backslash} q = \{p_0\}$ and $A \mathbin{\backslash\!\!\backslash} q' = \{p'\}$ with $p_0 = q$ or $p_0 = p'$.
- If $q' = p'q'_1$, $q'_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q'_1} \triangleleft l$, then $A \mathbin{\backslash\!\!\backslash} q' = \{p'q'_2 \mid q'_2 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(l) \wedge r|_{q'_1} = l|_{q'_2}\}$. Three cases from Definition 3.1 are applicable for q. Suppose $q < p'$, then $A \mathbin{\backslash\!\!\backslash} q = \{q\}$, and $q < p' \leqslant p'q'_2$ for any $q'_2 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(l)$. If $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q_1} \ntriangleleft l$, then $A \mathbin{\backslash\!\!\backslash} q = \{p'\}$. Last, if $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q_1} \triangleleft l$, then $A \mathbin{\backslash\!\!\backslash} q = \{p'q_2 \mid q_2 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(l) \wedge r|_{q_1} = l|_{q_2}\}$. We have $q' = qq'_3$, so for any $p'q_2 \in A \mathbin{\backslash\!\!\backslash} q$, also $p'q_2q'_3 \in A \mathbin{\backslash\!\!\backslash} q'$.
- Otherwise, $q' = p'q'_1q'_2$ with $q'_1 \in \mathcal{P}\mathsf{os}_{\mathcal{V}}(r)$. Then $A \mathbin{\backslash\!\!\backslash} q' = \{p'q'_3q'_2 \mid r|_{q'_1} = l|_{q'_3}\}$. Except for $q \parallel p'$, all cases in Definition 3.1 can happen for q. Suppose $q < p'$, then $A \mathbin{\backslash\!\!\backslash} q = \{q\}$, and $q < p' < p'q'_3q'_2$ for any $q'_3 \in \mathcal{P}\mathsf{os}_{\mathcal{V}}(l)$. If $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q_1} \ntriangleleft l$, then $A \mathbin{\backslash\!\!\backslash} q = \{p'\}$. For the next case, suppose $q = p'q_1$, $q_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$, and $r|_{q_1} \triangleleft l$. Then $A \mathbin{\backslash\!\!\backslash} q = \{p'q_2 \mid q_2 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(l) \wedge r|_{q_1} = l|_{q_2}\}$. We have $q' = qq'_4q'_2$, so for any $p'q_2 \in A \mathbin{\backslash\!\!\backslash} q$, also $p'q_2q'_4q'_2 \in A \mathbin{\backslash\!\!\backslash} q'$. Otherwise, $q = p'q'_1q_2$. Then $A \mathbin{\backslash\!\!\backslash} q = \{p'q_3q_2 \mid r|_{q'_1} = l|_{q_3}\}$. We have $q'_2 = q_2q'_4$, hence for any $p'q_3q_2 \in A \mathbin{\backslash\!\!\backslash} q$, also $p'q_3q'_2 \in A \mathbin{\backslash\!\!\backslash} q'$.

□

Notice that each position in a term may have several progenitors:

Example 3.3. Consider the TRS $\mathcal{R}_2$ consisting of the single rule $\mathsf{f}(x, x) \to g(x)$, and the rewrite step $A : \mathsf{f}(0, 0) \to_{\mathcal{R}_2} \mathsf{g}(0)$. Then $A \mathbin{\backslash\!\!\backslash} 1 = \{1, 2\}$.

We restrict the progenies and progenitors to a single branch in each term. The definition rests on the idea that for a derivation $A\colon s \to^* t$ and a main branch B' in t it is possible to find a *main branch* B in s such that each position $q \in B'$ has a (unique) progenitor in B; see the picture below for an illustration:



In the following definition, the restriction to the leftmost of all candidate positions is arbitrary and can be suitably replaced. Notice that its second clause is well-defined by Lemmata 3.3 and 3.4.

Definition 3.4. Let $A\colon t_1 \to^* t_n$ denote a derivation built up from the rewrite steps $A_i\colon t_i \to t_{i+1}$ for $i = 1, \ldots, n-1$. Then the *main branch* of each term in A is inductively defined:

- The main branch of t_n is the leftmost branch of maximal length in t_n.
- Suppose the main branch of t_{i+1} is denoted as B_{i+1}, $1 \leqslant i \leqslant n-1$. Then consider all branches b in t_i such that for every $q \in B_{i+1}$, the set of progenitors $A_i \mathbin{\backslash\backslash} q$ of q has nonempty intersection with b. The leftmost of these branches is the main branch of t_i, denoted as B_i.

Definition 3.5. Let $A' : s \to t$ be a rewriting step, let $p \in \mathcal{P}\mathsf{os}(s)$, and let B and B' be branches in s and t. Then the set of *main progenies of p in t* (with respect to A') (denoted as $p \mathbin{\text{Ə}}^{B}_{B'} A'$) is defined as follows:

$$p \mathbin{\text{Ə}}^{B}_{B'} A' = \begin{cases} \varnothing & \text{if } p \notin B \\ B' \cap p \mathbin{\backslash\backslash} A' & \text{if } p \in B \end{cases}$$

We naturally extend this definition to derivations $A : s \to^* t$, analogous to Definition 3.2: If A is the empty derivation, then $p \mathbin{\text{Ə}}^{B}_{B'} A = \{p\}$. Otherwise, we can split A into $A_1 : s \to s'$ and $A_2 : s' \to^* t$. Let B'' be the main branch in s'. Then $p \mathbin{\text{Ə}}^{B}_{B'} A = (p \mathbin{\text{Ə}}^{B}_{B''} A_1) \mathbin{\text{Ə}}^{B''}_{B'} A_2$. If the (main) branches B and B' are clear from context, we write $p \mathbin{\text{Ə}} A'$ instead of $p \mathbin{\text{Ə}}^{B}_{B'} A'$. If $q \in p \mathbin{\text{Ə}} A'$, then we also say that p is a *main progenitor of q in s* (with respect to A'). We denote the set of main progenitors of q in s by $A' \mathbin{\text{Ə}} q$. For a set $P \subseteq \mathcal{P}\mathsf{os}(s')$, we define $P \mathbin{\text{Ə}} A' = \bigcup_{p \in P} p \mathbin{\text{Ə}} A'$.

Remark 3.1. Note that it is indeed necessary to define main progenies with respect to many-step derivations in the inductive way shown above. Consider the TRS with the rewrite rules $\mathsf{f}(x) \to \mathsf{g}(x, \mathsf{c})$ and $\mathsf{g}(x, x) \to \mathsf{h}(x, x)$, and the derivation $A : t_1 \to t_2 \to t_3$, where $t_1 = \mathsf{f}(\mathsf{c})$, $t_2 = \mathsf{g}(\mathsf{c}, \mathsf{c})$, and $t_3 = \mathsf{h}(\mathsf{c}, \mathsf{c})$. We split A into $A_1 : t_1 \to t_2$ and $A_2 : t_2 \to t_3$. The main branch of each term in A is the leftmost one (i.e. $\{\epsilon, 1\}$). We obviously have $1 \in A \mathbin{\text{Ə}} 1$, since $1 \in A_2 \mathbin{\text{Ə}} 1$ and $1 \in A_1 \mathbin{\text{Ə}} 1$. However, we do not have $\epsilon \in A \mathbin{\text{Ə}} 1$, even though $2 \in A_2 \mathbin{\backslash\backslash} 1$, $\epsilon \in A_1 \mathbin{\backslash\backslash} 2$, and therefore $\epsilon \in A \mathbin{\backslash\backslash} 1$. Hence, the direct definition of main progenies with respect to many-step derivations is indeed different from the definition we are using. In particular, Lemma 3.5 would be incorrect for that definition.

Example 3.4 (continued from Example 3.1)**.** Consider the derivation A from Example 3.1. The "central" branch of each term in Figure 1 is its main branch, and the dashed lines denote the main progeny relation.

Lemma 3.5. *Let $A\colon u \to^* s \to^* t \to^* w$ be a derivation, and denote $A'\colon s \to^* t$. Let $B(s)$ and $B(t)$ denote the main branches of s and t, respectively. Then for any $q \in B(t)$, the main progenitor of q in branch $B(s)$ is unique, i.e., $|A' \mathbin{\text{Ə}} q| = 1$.*

Proof. By definition, q has at least one main progenitor in s. We show that there exists at most one by induction on the length n of $A'\colon s \to^n t$. For $n = 0$ the claim is trivial. Hence assume $n > 0$ and let $A'\colon s \to t' \to^{n-1} t$. Let $B(t')$ denote the main branch in t' (with respect to A). By induction hypothesis there exists a unique position q_1 in $B(t')$ such that $(t' \to^{n-1} t) \mathbin{\text{Ə}} q = \{q_1\}$. Let $A''\colon s \to_{p', l \to r} t'$ denote the first rewrite step in A'. Suppose $q_1 < p'$ or $q_1 \parallel p'$. Then by definition $A'' \mathbin{\backslash\!\backslash} q_1 = \{q_1\}$. Hence the main progenitor of q in $B(s)$ is unique. On the other hand suppose $q_1 = p'q_2$ with $q_2 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$ such that $r|_{q_2} \ntriangleleft l$. Then $A'' \mathbin{\backslash\!\backslash} q_1 = \{p'\}$ and $A' \partial q$ is a singleton as it should be. Now suppose $q_1 = p'q_2$ with $q_2 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(r)$ such that $r|_{q_2} \triangleleft l$. Then by definition $A'' \mathbin{\backslash\!\backslash} q_1 = \{p'p_1 \mid p_1 \in \mathcal{P}\mathsf{os}_{\mathcal{F}}(l) \wedge l|_{p_1} = r|_{q_2}\}$. Notice that $A' \partial q = A'' \mathbin{\backslash\!\backslash} q_1 \cap B(s)$, which is again a singleton. Finally, if $q_1 = p'q_2q_3$ with $q_2 \in \mathcal{P}\mathsf{os}_{\mathcal{V}}(r)$, then $A'' \mathbin{\backslash\!\backslash} q_1 = \{p'p_1q_3 \mid p_1 \in \mathcal{P}\mathsf{os}_{\mathcal{V}}(l) \wedge l|_{p_1} = r|_{q_2}\}$. As before, the intersection of the latter set with $B(s)$ is a singleton. Hence the main progenitor of q in $B(s)$ is unique. This concludes the inductive proof. □

Lemma 3.6. *We assume the same notation as in Lemma 3.5. For any* $p \in B(s)$ *such that* $\mathsf{rt}(s|_p) \in \mathcal{C} \cup \mathcal{V}$*, we have* $|p \partial A'| \leqslant 1$*, i.e., the number of main progenies for a position, whose root is non-defined is at most* 1.

Proof. By induction on the length n of $A'\colon s \to^n t$. It suffices to consider the case $n > 0$, so $A'\colon s \to t' \to^{n-1} t$. Let $A''\colon s \to_{p',l\to r} t'$ denote the first rewrite step in A'. If $p < p'$ or $p \parallel p'$, then $p \partial A'' = \{p\}$. If $p > p'$, then for any $q_1 \in p \mathbin{\backslash\!\backslash} A''$, we have $s|_p = t'|_{q_1}$, so again, $p \mathbin{\backslash\!\backslash} A'' \cap B(t')$ is a singleton. In all three cases, the claim follows by induction hypothesis as $\mathsf{rt}(s|_p) = \mathsf{rt}(t|_{q_1})$. This concludes the proof, as the case $p = p'$ is impossible. Otherwise, we derive a contradiction to the assumption that the root of $s|_p$ is not a defined symbol. □

4 Dependency Pairs and Complexity

In this section, we relate the dependency pair complexity and the derivational complexity of a TRS. As mentioned above the main step is to show that the depth growth of a TRS is bounded by a single exponential in its dependency pair complexity. In the sequel, we fix a TRS $\mathcal{R}$ and a derivation $A : t_1 \to^* t_n$ over $\mathcal{R}$ such that $B_1, \ldots, B_n$ denote the main branches with respect to A. We can view the main progeny relation as a graph, called *progenitor graph*. The nodes of a progenitor graph are pairs (t_i, p) representing positions p in terms in A that are directly affected by dependency pair steps. Each edge corresponds to a dependency pair step (and possibly a number of $\mathcal{R}$-steps) in an implicit dependency pair derivation. Thus we obtain a graph representation of all implicit dependency pair derivations in A. Each connected component of the progenitor graph is a tree whose height is bounded by the number of dependency pair steps. Using the exponential relationship between the height of this tree and its number of leaves, we bound the depth of the final term in A exponentially in the length of the largest implicit dependency pair derivation, entailing our main result.

Definition 4.1. The *progenitor graph* of A is defined as follows.

- The nodes are all pairs (t_i, p) such that $p \in B_i$, $\mathsf{rt}(t_i|_p) \in \mathcal{D}$, and either $i = 1$ or the single element of $(t_{i-1} \to t_i) \partial p$ and the redex position in the rewrite step $t_{i-1} \to t_i$ coincide.

- There is an edge from (t_i, p) to (t_j, q) whenever $i < j$, $(t_i \to^* t_j) \partial q = \{p\}$, and for all $i \leqslant k < j-1$, the single element of $(t_k \to^* t_j) \partial q$ and the redex position in the rewrite step $t_k \to t_{k+1}$ do not coincide.

Notice that, due to the definition of the set of nodes in a progenitor graph, the single element of $(t_k \to^* t_j) \partial q$ and the redex position in the rewrite step $t_{j-1} \to t_j$ do coincide whenever there is an edge to (t_j, q).

Example 4.1. Consider the derivation A from Example 3.1 again. Its progenitor graph is shown below:



Lemma 4.1. *If there is an edge from (t_i, p) to (t_j, q) in G, then there is a derivation $(t_i|_p)^\sharp \to^*_{\mathcal{R}} (t_{j-1}|_{p_1})^\sharp \to_{\mathsf{DP}(\mathcal{R})} (t_j|_q)^\sharp$.*

Proof. By definition, $q \in p \partial (t_i \to^* t_j)$. Therefore, by Lemma 3.3, we have the implicit dependency pair derivation $A' : (t_i|_p)^\sharp \to^*_{\mathcal{R} \cup \mathsf{DP}(\mathcal{R})} (t_j|_q)^\sharp$. We have $(t_{j-1} \to t_j) \partial q = \{p_1\}$ (see also the last clause of Definition 3.1), where by definition p_1 is the redex position of the step $t_{j-1} \to t_j$. Therefore, the last step of A' is a $\mathsf{DP}(\mathcal{R})$-step. Notice that for $i \leqslant k < j-1$, the single element of $(t_k \to^* t_j) \partial q$ and the redex position in $t_k \to t_{k+1}$ do not coincide. Hence, if there are rewrite steps before the last step, these are $\mathcal{R}$-steps and the lemma follows. ☐

The next lemma shows, when specialised to the conditions in the first clause of Definition 4.1, that only nodes which do not contribute to the branching of the progenitor graph, are left out by the definition.

Lemma 4.2. *Let $p \in B_i$ and $q \in B_j$ such that $i < j$ and $(t_i \to^* t_j) \partial q = \{p\}$. If for all $i \leqslant k \leqslant j-1$, the single element of $(t_k \to^* t_j) \partial q$ and the redex position in the rewrite step $t_k \to t_{k+1}$ do not coincide, then $p \partial (t_i \to^* t_j) = \{q\}$.*

Proof. We show the lemma by induction on $j - i$. If $i = j$ then the claim trivially holds. Otherwise, the derivation $t_i \to^* t_j$ can be split to $t_i \to t_{i+1} \to^* t_j$. Let p' be the redex position in $t_i \to t_{i+1}$. If $p \parallel p'$, $p < p'$, or $p > p'$, then as in Lemma 3.6, $|p \partial (t_i \to t_{i+1})| \leqslant 1$, and the lemma follows by induction hypothesis. The last case is again impossible, since by assumption, p and p' do not coincide. ☐

In the sequel let G be the progenitor graph of A. In the next lemmata, we show the properties which allow us to bound $\mathsf{dp}(t_n)$ in the height of G. First, we prove that almost each position in B_n is "covered" by a node in G. Next, we show that each node in G can only cover c positions in B_n, and finally, we show that the branching factor of G is at most c, where $c := \max\{2\} \cup \{\mathsf{dp}(r) \mid l \to r \in \mathcal{R}\}$.

Lemma 4.3. *For every* $q \in B_n$*, there either exists* $p \in B_1$ *such that* $\mathsf{rt}(t_1|_p) \in \mathcal{C} \cup \mathcal{V}$ *and* $A \eth q = \{p\}$*, or there exists a node* (t_i, p) *in* G *where* $q \in p \eth (t_i \to^* t_n)$ *and for any successor node* (t_j, p_1) *of* (t_i, p) *in* G*, we have* $q \notin p_1 \eth (t_j \to^* t_n)$*.*

Proof. By Lemma 3.5, $A \eth q = \{p\}$ for some $p \in B_1$. If $\mathsf{rt}(t_1|_p) \in \mathcal{C} \cup \mathcal{V}$, the first alternative of the lemma holds. If $\mathsf{rt}(t_1|_p) \in \mathcal{D}$, then $(t_1, p) \in G$. Therefore, there exists a maximal natural number k such that $(t_k, p_2) \in G$ and $q \in p_2 \eth (t_k \to^* t_n)$ for some $p_2 \in B_k$, so the second alternative of the lemma holds for (t_k, p_2). ☐

Lemma 4.4. *For every node* (t_i, p) *in* G*, there are at most* c *many positions* $q \in B_n$ *such that* $q \in p \eth (t_i \to^* t_n)$*, but for any successor node* (t_j, p_1) *of* (t_i, p)*, we have* $q \notin p_1 \eth (t_j \to^* t_n)$*.*

Proof. If there is no $i \leqslant k < n$ such that the redex position of the step $t_k \to t_{k+1}$ and an element of $p \eth (t_i \to^* t_k)$ coincide, then it follows from Lemma 4.2 that $|p \eth (t_i \to^* t_n)| \leqslant 1$. Otherwise, let k be the smallest number such that $k \geqslant i$ and $p \eth (t_i \to^* t_k) = \{p_2\}$, where p_2 is the redex position of $t_k \to t_{k+1}$. By Definitions 3.1 and 3.5, $|p_2 \eth (t_k \to t_{k+1})| \leqslant c$. For each $p_3 \in p_2 \eth (t_k \to t_{k+1})$, if $\mathsf{rt}(t_{k+1}|_{p_3}) \in \mathcal{D}$, then (t_{k+1}, p_3) is a successor node of (t_i, p), but the condition $q \notin p_3 \eth (t_{k+1} \to^* t_n)$ is violated for any main progeny q of p_3. On the other hand, if $\mathsf{rt}(t_{k+1}|_{p_3}) \in \mathcal{C} \cup \mathcal{V}$, then by Lemma 3.6, $|p_3 \eth (t_{k+1} \to^* t_n)| \leqslant 1$. Thus, in total, there are at most c many elements in $p \eth (t_i \to^* t_n)$ meeting our assumption. ☐

The following example illustrates the role of Lemma 4.4.

Example 4.2. Let $\mathcal{R}_3$ be the TRS consisting of the single rewrite rule

$$\mathsf{d}(\mathsf{S}(x)) \to \mathsf{S}(\mathsf{S}(\mathsf{d}(x))) \ .$$

Let $t_1 = \mathsf{d}(\mathsf{S}(\mathsf{S}(0)))$, $t_2 = \mathsf{S}(\mathsf{S}(\mathsf{d}(\mathsf{S}(0))))$, and $t_3 = \mathsf{S}(\mathsf{S}(\mathsf{S}(\mathsf{S}(\mathsf{d}(0)))))$. We have the derivation $A : t_1 \to t_2 \to t_3$ and the following progenitor graph:



Note that G leaves out all function symbols S above the d in each term. However, by Lemma 4.4, the number of positions in the last term of A which are hidden in this way is only linear in the size of the progenitor graph.

Lemma 4.5. *Every node in* G *has at most* c *many successor nodes.*

Proof. Let (t_i, p) be a node in G. If there is no $i \leqslant j < n$ such that the redex position of the step $t_j \to t_{j+1}$ and an element of $p \eth (t_i \to^* t_j)$ coincide, then (t_i, p) has no successor node, so the claim holds. Otherwise, let j be the smallest number greater than i such that $p \eth (t_i \to^* t_j) = \{q\}$, where q is the redex position of $t_j \to t_{j+1}$. By Definitions 3.1 and 3.5, $|q \eth (t_j \to t_{j+1})| \leqslant c$. Hence, (t_i, p) has at most c many successor nodes. ☐

Now we are ready to prove our main lemma.

Lemma 4.6. *For every finite TRS $\mathcal{R}$, there exists a constant C such that for all terminating terms s, we have* $\mathsf{pdp}(s, \to_{\mathcal{R}}) \leqslant |s| \cdot 2^{C \cdot \mathsf{dl}(s^\sharp, \to_{\mathsf{DP}(\mathcal{R})/\mathcal{R}})}$.

Proof. We show the lemma by proving that for any derivation $A : s \to_{\mathcal{R}}^* t$, there exists a derivation $A' : (s')^\sharp \to_{\mathcal{R} \cup \mathsf{DP}(\mathcal{R})}^* (t')^\sharp$ such that $|s'| \leqslant |s|$ and $\mathsf{dp}(t) \leqslant |s| \cdot c^{\mathsf{DPl}(A')+2}$ (recall that $c = \max\{2\} \cup \{\mathsf{dp}(r) \mid l \to r \in \mathcal{R}\}$). Let k be the number of defined symbols in the main branch of s. Observe that the main branch of t consists of $\mathsf{dp}(t) + 1$ many positions, all of which have to fulfil one of the two properties outlined in Lemma 4.3. By Lemma 3.6, the first case applies to at most $\mathsf{dp}(s) + 1 - k$ many positions, so for the $\mathsf{dp}(t) - \mathsf{dp}(s) + k$ other positions, the second case applies. By Lemma 4.4, each node in the progenitor graph G of A can cover at most c many of those positions, so G has to contain at least $\frac{\mathsf{dp}(t) - \mathsf{dp}(s) + k}{c}$ many nodes. There are k many connected components in G, hence the largest one of the components (trees) contains at least

$$\frac{\mathsf{dp}(t) - \mathsf{dp}(s) + k}{kc} ,$$

many nodes. Let $d - 2$ be the smallest natural number such that

$$\frac{\mathsf{dp}(t) - \mathsf{dp}(s) + k}{kc} \leqslant c^{d-1} .$$

By Lemma 4.5, this means that there exists a leaf whose distance from the root is at least d. Recall that any c-ary tree of height $d - 1$ has at most $\frac{c^{d-1} - 1}{c - 1} \leqslant c^{d-1}$ many nodes. (Here the height of a tree is the number of nodes from the root to the leaf on a maximal path.) Moreover, by Lemma 4.1, this path in the graph induces a derivation $A' : (s')^\sharp \to_{\mathcal{R} \cup \mathsf{DP}(\mathcal{R})}^* (t')^\sharp$ with $\mathsf{DPl}(A') \geqslant d - 2$ and $s \trianglerighteq s'$. Reformulating the inequality above yields

$$\mathsf{dp}(t) \leqslant k \cdot c^d + \mathsf{dp}(s) - k \leqslant (\mathsf{dp}(s) + 1) \cdot c^d \leqslant |s| \cdot c^d ,$$

so A' is indeed the derivation we are looking for. ☐

The main factor of the faster growth of $\mathsf{dp}(t_n)$ compared to the height of G is the difference between the height and the size of G. This becomes apparent in our next example, where G is a full binary tree.

Example 4.3. Consider the TRS $\mathcal{R}_4$ consisting of the single rewrite rule

$$\mathsf{f}(\mathsf{S}(x), y) \to \mathsf{f}(x, \mathsf{f}(x, y)) .$$

Consider the terms $t_1 = \mathsf{f}(\mathsf{S}(\mathsf{S}(\mathsf{0})), \mathsf{0})$, $t_2 = \mathsf{f}(\mathsf{S}(\mathsf{0}), \mathsf{f}(\mathsf{S}(\mathsf{0}), \mathsf{0}))$, $t_3 = \mathsf{f}(\mathsf{0}, \mathsf{f}(\mathsf{0}, \mathsf{f}(\mathsf{S}(\mathsf{0}), \mathsf{0})))$, and $t_4 = \mathsf{f}(\mathsf{0}, \mathsf{f}(\mathsf{0}, \mathsf{f}(\mathsf{0}, \mathsf{f}(\mathsf{0}, \mathsf{0}))))$. We have the derivation $A : t_1 \to t_2 \to t_3 \to t_4$. The progenitor graph of A is shown below.



Perhaps counter-intuitively in the context of the dependency pair method, the connected component of G with the greatest height need not be the component with the root (t_1, ϵ):

Example 4.4. Consider the TRS $\mathcal{R}_4$ and the terms t_1, t_2, t_3, t_4 from Example 4.3 again. Let $s_1 = \mathsf{f}(\mathsf{0}, t_1)$, $s_2 = \mathsf{f}(\mathsf{0}, t_2)$, $s_3 = \mathsf{f}(\mathsf{0}, t_3)$, and $s_4 = \mathsf{f}(\mathsf{0}, t_4)$. Then we obviously have the derivation $A' : s_1 \to s_2 \to s_3 \to s_4$ again and the following progenitor graph:



All that is left to show is that the derivational complexity of a finite and terminating TRS is bounded double exponentially in its depth growth. This can be achieved by two easy observations.

Lemma 4.7. *Let $\mathcal{R}$ be a finite TRS. Then there exists a constant C such that for every terminating term t, we have*

$$\mathsf{dl}(t, \to_{\mathcal{R}}) \leqslant 2^{2^{C \cdot \mathsf{pdp}(t, \to_{\mathcal{R}})}}$$

Proof. We show that there exist constants D and E, such that for all terms t, the inequalities $\mathsf{psz}(t, \to_{\mathcal{R}}) \leqslant 2^{D \cdot \mathsf{pdp}(t, \to_{\mathcal{R}})}$ and $\mathsf{dl}(t, \to_{\mathcal{R}}) \leqslant 2^{E \cdot \mathsf{psz}(t, \to_{\mathcal{R}})}$ hold.

a. For any term t, we have $|t| \leqslant k^{\mathsf{dp}(t)+1}$, where k is the maximum arity of any function symbol in the signature. This proves the first inequality.

b. On the other hand, by assumption the signature $\mathcal{F}$ of $\mathcal{R}$ is finite. Moreover without loss of generality the considered derivation in $\mathcal{R}$ is ground. Hence we can build only

$2^{E \cdot m}$ different terms of size at most m, where E depends only on $\mathcal{F}$. This proves the second inequality.

$\square$

Based on Lemmata 4.6 and 4.7 we obtain our main theorem.

Theorem 4.1. *For any finite and terminating TRS* $\mathcal{R}$, $\mathsf{dc}_{\mathcal{R}}(n) \leqslant 2^{2^{n \cdot 2^{O(\mathsf{DPc}_{\mathcal{R}}(n))}}}$.

An order $\succ$ on terms is G-*collapsible* for a TRS $\mathcal{R}$ if $s \rightarrow_{\mathcal{R} \cup \mathsf{DP}(\mathcal{R})} t$ and $s \succ t$ implies $\mathsf{G}(s, \succ) > \mathsf{G}(t, \succ)$. Let $(\gtrsim, \succ)$ be a reduction pair for $\mathcal{R}$. Then $(\gtrsim, \succ)$ is called *collapsible* if there is a mapping G such that $\succ$ is G-collapsible for $\mathcal{R}$.

Theorem 4.2. *Let* $\mathcal{R}$ *be a finite TRS, let* $(\gtrsim, \succ)$ *be a collapsible reduction pair with* $\pi(\mathcal{R}) \subseteq \gtrsim$ *and* $\pi(\mathsf{DP}(\mathcal{R})) \subseteq \succ$ *for some argument filtering* π. *Assume there exists a class of number-theoretic functions* $\mathcal{C}$ *closed under elementary functions and for some* $f \in \mathcal{C}$, *and any term* t, $\mathsf{G}(t, \succ) \leqslant f(|t|)$. *Then* $\mathsf{dc}_{\mathcal{R}} \in \mathcal{C}$.

Proof. By assumption there exists a mapping G that binds the number of dependency pair steps in any $\pi(\mathcal{R}) \cup \pi(\mathsf{DP}(\mathcal{R}))$-derivation. More precisely, we have

$$\mathsf{dl}(\pi(t^\sharp), \rightarrow_{\pi(\mathsf{DP}(\mathcal{R}))/\pi(\mathcal{R})}) \leqslant \mathsf{G}(t, \succ) \leqslant f(|t|) \; . \tag{1}$$

Moreover it is easy to see that for any derivation in $\mathcal{R} \cup \mathsf{DP}(\mathcal{R})$, there is a derivation in $\pi(\mathcal{R}) \cup \pi(\mathsf{DP}(\mathcal{R}))$ which contains the same number of dependency pair steps. Hence, we obtain

$$\mathsf{dl}(t^\sharp, \rightarrow_{\mathsf{DP}(\mathcal{R})/\mathcal{R}}) \leqslant \mathsf{dl}(\pi(t^\sharp), \rightarrow_{\pi(\mathsf{DP}(\mathcal{R}))/\pi(\mathcal{R})}) \; .$$

Combining this with (1) and Theorem 4.1 we obtain $\mathsf{dc}_{\mathcal{R}}(n) \leqslant 2^{2^{n \cdot 2^{a \cdot f(n)}}}$. By assumption the complexity class $\mathcal{C}$ is closed under elementary functions. In particular there exists $g \in \mathcal{C}$ such that $\mathsf{dc}_{\mathcal{R}}(n) \leqslant g(n)$. Thus the theorem follows. $\square$

5 The Lower Bound

By Theorem 4.1, the derivational complexity of a TRS $\mathcal{R}$ is bounded triple exponentially in its dependency pair complexity. This yields an upper bound. The following TRS establishes a double exponential lower bound.

Example 5.1. Consider the following TRS $\mathcal{R}_5$, extending the TRS $\mathcal{R}_4$.

$$1\colon\; \mathsf{f}(\mathsf{S}(x), y) \rightarrow \mathsf{f}(x, \mathsf{f}(x, y)) \qquad 2\colon\; \mathsf{f}(\mathsf{0}, x) \rightarrow \mathsf{c}(x, x)$$

We show that $\mathcal{R}_5$ has linear dependency pair complexity, but admits derivations of double exponential length. Let $F^0_m(x) = x$, $F^{n+1}_m(x) = \mathsf{f}(\mathsf{S}^m(\mathsf{0}), F^n_m(x))$, $C^0(x) = x$, and $C^{n+1}(x) = \mathsf{c}(C^n(x), C^n(x))$. Now, consider the starting term $F^1_n(\mathsf{0})$. As can be easily seen, this term rewrites to $F^{2^n}_0(\mathsf{0})$ in $2^n - 1$ steps using rule 1. Now, we can use rule 2 and an outermost strategy to reach $C^{2^n}(0)$ in $2^{2^n}-1$ steps, so $\mathsf{dc}_{\mathcal{R}_5}(n)$ is at least double exponential in n. On the other hand consider $\mathsf{DP}(\mathcal{R}_5)$:

$$3\colon\ \mathsf{f}^\sharp(\mathsf{S}(x),y) \to \mathsf{f}^\sharp(x,\mathsf{f}(x,y)) \qquad 4\colon\ \mathsf{f}^\sharp(\mathsf{S}(x),y) \to \mathsf{f}^\sharp(x,y)$$

We define a (restricted) polynomial interpretation $\mathcal{A}$ as follows: $\mathsf{f}^\sharp_{\mathcal{A}}(x,y) = x$, $\mathsf{S}_{\mathcal{A}}(x) = x+1$, $\mathsf{f}_{\mathcal{A}}(x,y) = \mathsf{c}_{\mathcal{A}}(x,y) = \mathsf{0}_{\mathcal{A}} = 0$, such that $\mathcal{R}_5 \subseteq {\gtrsim_{\mathcal{A}}}$ and $\mathsf{DP}(\mathcal{R}_5) \subseteq {>_{\mathcal{A}}}$, and $(\gtrsim_{\mathcal{A}}, >_{\mathcal{A}})$ forms a reduction pair. Thus the dependency pair complexity of $\mathcal{R}_5$ is at most linear.

Notice that from the proof of Theorem 4.1 one can distill the following three facts, where each of them seems to be responsible for one of the exponentials in the upper bound:

- The number of leaves in a progenitor graph may be exponential in its height.
- The size of a term may be exponential in its depth.
- The number of terms of size n is exponential in n.

Observe that for an optimal example, we would have to utilise all three criteria, while the just given TRS $\mathcal{R}_5$ utilises only the first two criteria. At this point, it seems impossible to enumerate enough terms of exponential depth and double exponential size so that this is possible. Hence, we conjecture that the upper bound given in Theorem 4.1 can be improved to double exponential.

6 Dependency Graphs

We now consider *dependency graphs*, which are another improvement of the dependency pair method besides argument filtering. At first glance, it might seem that the number of dependency pair steps admitted by a TRS is bounded linearly in the number of dependency pair steps admitted within the “worst” SCC of the dependency graph. However, this is not the case:

Example 6.1. Consider the following TRS $\mathcal{R}_{\mathsf{e}_2}$:

$$\begin{array}{lll} \mathsf{d}(\mathsf{0}) \to \mathsf{0} & \mathsf{e}_2(\mathsf{0},x) \to \mathsf{e}_1(x,\mathsf{S}(\mathsf{0})) & \mathsf{e}_1(\mathsf{S}(x),y) \to \mathsf{e}_1(x,\mathsf{d}(y)) \\ \mathsf{d}(\mathsf{S}(x)) \to \mathsf{S}(\mathsf{S}(\mathsf{d}(x))) & \mathsf{e}_2(\mathsf{S}(x),y) \to \mathsf{e}_2(x,\mathsf{d}(y)) & \end{array}$$

We have the following set of dependency pairs $\mathsf{DP}(\mathcal{R}_{\mathsf{e}_2})$:

$$\begin{array}{lrlr} \mathsf{d}^\sharp(\mathsf{S}(x)) \to \mathsf{d}^\sharp(x) & (1) & \mathsf{e}_2^\sharp(\mathsf{0},x) \to \mathsf{e}_1^\sharp(x,\mathsf{S}(\mathsf{0})) & (4) \\ \mathsf{e}_1^\sharp(\mathsf{S}(x),y) \to \mathsf{e}_1^\sharp(x,\mathsf{d}(y)) & (2) & \mathsf{e}_2^\sharp(\mathsf{S}(x),y) \to \mathsf{e}_2^\sharp(x,\mathsf{d}(y)) & (5) \\ \mathsf{e}_1^\sharp(\mathsf{S}(x),y) \to \mathsf{d}^\sharp(y) & (3) & \mathsf{e}_2^\sharp(\mathsf{S}(x),y) \to \mathsf{d}^\sharp(y) & (6) \end{array}$$

This induces the dependency graph $\mathsf{DG}(\mathcal{R}_{\mathsf{e}_2})$ shown in Figure 2. The dependency graph

Figure 2: Dependency Graph for the TRS $\mathcal{R}_{\mathsf{e}_2}$



contains three (nontrivial) SCCs: $\{1\}$, $\{2\}$, and $\{5\}$. For each of the SCCs, we can prove termination (and an exponential upper bound on the number of dependency pair steps within that SCC) by a linear polynomial algebra $\mathcal{A}$ with

$$\mathsf{d}_{\mathcal{A}}(x) = 2x \ , \quad \mathsf{e}_{1\mathcal{A}}(x,y) = 0 \ , \quad \mathsf{e}_{2\mathcal{A}}(x,y) = 0 \ , \quad \mathsf{S}_{\mathcal{A}}(x) = x+1 \ , \quad \mathsf{0}_{\mathcal{A}} = 0 \ .$$

For the SCC $\{1\}$, we can extend $\mathcal{A}$ with $\mathsf{d}^{\sharp}_{\mathcal{A}}(x) = x+1$, for $\{2\}$, we can set $e^{\sharp}_{1\mathcal{A}}(x,y) = x$, and for the SCC $\{5\}$, we have $e^{\sharp}_{2\mathcal{A}}(x,y) = x$.

However, the full dependency pair complexity admitted by $\mathcal{R}_{\mathsf{e}_2}$ is at least double exponential: we have $\mathsf{e}^{\sharp}_2(\mathsf{S}^n(\mathsf{0}),\mathsf{S}(\mathsf{0})) \to^* \mathsf{e}^{\sharp}_1(\mathsf{S}(\mathsf{0}),\mathsf{S}^{2^n}(\mathsf{0})) \to^* \mathsf{d}^{\sharp}(\mathsf{S}^{2^{2^n}-1}(\mathsf{0})) \to_{\mathsf{DP}(\mathcal{R}_{\mathsf{e}_2})}^{2^{2^n}-1} \mathsf{d}^{\sharp}(\mathsf{0})$.

Remark 6.1. Note that one may be tempted to incorrectly conclude that the number of dependency pair steps in each SCC is linear by the subterm criterion. Indeed, each SCC in this example can be proved terminating by the subterm criterion. However, contrary to what one may think, the subterm criterion does not give a linear complexity bound. While the derivational complexity of the embedding rules is obviously linear, the rewrite rules applied between the dependency pair steps may increase the size of the given term arbitrarily.

Example 6.2. We can generalise the TRS $\mathcal{R}_{\mathsf{e}_2}$ to TRSs $\mathcal{R}_{\mathsf{e}_k}$ for $k \in \mathbb{N} \setminus \{0\}$, such that each cycle in $\mathsf{DG}(\mathcal{R}_{\mathsf{e}_k})$ admits only exponentially many dependency pair steps, but whose full dependency pair complexity is $2_k(\Theta(n))$. The rules of $\mathcal{R}_{\mathsf{e}_k}$ are the following:

$$\begin{aligned}
\mathsf{d}(\mathsf{0}) &\to \mathsf{0} \\
\mathsf{d}(\mathsf{S}(x)) &\to \mathsf{S}(\mathsf{S}(\mathsf{d}(x))) \\
\mathsf{e}_i(\mathsf{S}(x),y) &\to \mathsf{e}_i(x,\mathsf{d}(y)) && \text{whenever } 1 \leqslant i \leqslant k \\
\mathsf{e}_i(\mathsf{0},x) &\to \mathsf{e}_{i-1}(x,\mathsf{S}(\mathsf{0})) && \text{whenever } 2 \leqslant i \leqslant k
\end{aligned}$$

The exponential upper bound in the number of dependency pair steps in each SCC of $\mathsf{DG}(\mathcal{R}_{\mathsf{e}_k})$ can be shown in a similar way as in Example 6.1. Moreover, we can also verify that the dependency pair complexity of $\mathcal{R}_{\mathsf{e}_k}$ is at least $2_k(\Theta(n))$ by looking at the longest derivation started from $\mathsf{e}^{\sharp}_k(\mathsf{S}^n(\mathsf{0}),\mathsf{S}(\mathsf{0}))$.

The problem in Examples 6.1 and 6.2 is that the bounds on the dependency pair complexity of the SCCs are related to the size of the considered term at the moment of entering the SCC, and not to the size of the starting term of the full derivation. Thus, Theorem 4.1 can not be extended directly. However, the maximal number of dependency pair steps allowed in the SCCs is related to the dependency pair complexity of the whole TRS as follows:

Theorem 6.1. *Let $\mathcal{R}$ be a finite TRS, let k be the number of (trivial and nontrivial) SCCs in a sound approximation of $\mathsf{DG}(\mathcal{R})$, and let f be a function bounding $\mathsf{dl}(u^\sharp, \to_{\mathcal{C}/\mathcal{R}})$ in $|u^\sharp|$ for each SCC $\mathcal{C}$. Then for any term t, $\mathsf{dl}(t^\sharp, \to_{\mathsf{DP}(\mathcal{R})/\mathcal{R}}) \leqslant 2^{F^k(|t^\sharp|)}$, where $F(n) = 2^{n \cdot 2^{a \cdot f(n)}}$ for some constant $a > 0$.*

Proof. In any derivation, whenever a dependency pair step is done, then the next dependency pair step in that chain may only stem from a dependency pair that is a successor (in $\mathsf{DG}(\mathcal{R})$) of the dependency pair applied in the current step. Therefore, the derivation can be split into $l \leqslant k$ segments such that in each of these segments, only dependency pairs from a single one of the SCCs in the approximation of $\mathsf{DG}(\mathcal{R})$ are applied in dependency pair steps. We now show $\mathsf{dl}(t^\sharp, \to_{\mathsf{DP}(\mathcal{R})/\mathcal{R}}) \leqslant 2^{F^l(|t^\sharp|)}$ by induction on l. If $l = 1$, then the claim follows directly from Theorem 4.1. Otherwise, consider the first segment of the derivation, let $t^\sharp$ be its starting term, and let $u^\sharp$ be the starting term of the second segment. By Lemmata 4.6 and 4.7, we have $|u^\sharp| \leqslant F(|t^\sharp|)$. By induction hypothesis, $\mathsf{dl}(u^\sharp, \to_{\mathsf{DP}(\mathcal{R})/\mathcal{R}}) \leqslant 2^{F^{l-1}(|u^\sharp|)}$. Putting the two inequalities together, we get the claim, and thus conclude the inductive proof. $\square$

Note that if $f(n)$ is an elementary, primitive recursive, or multiply recursive function, then $2^{F^k(n)}$ is still an elementary, primitive recursive, or multiply recursive function, respectively. The next theorem is a direct consequence of Theorems 4.2 and 6.1.

Theorem 6.2. *Let $\mathcal{R}$ be a finite TRS, and let k be the number of (trivial and nontrivial) SCCs in a sound approximation of $\mathsf{DG}(\mathcal{R})$. Assume there exists a class of number-theoretic functions $\mathcal{C}$ closed under elementary functions, and for some $f \in \mathcal{C}$, any term u, and any SCC $\mathcal{C}$, $\mathsf{G}(\pi(u^\sharp), \succ) \leqslant f(|u|)$, where $(\gtrsim, \succ)$ is a collapsible reduction pair with $\pi(\mathcal{R}) \subseteq \gtrsim$ and $\pi(\mathcal{C}) \subseteq \succ$ for some argument filtering π. Then $\mathsf{dc}_{\mathcal{R}} \in \mathcal{C}$.*

7 Conclusion

In this paper we have shown that the derivational complexity of a TRS $\mathcal{R}$ is bounded triple exponentially in its dependency pair complexity. Moreover we have presented an example showing that the relationship is at least double exponential. Furthermore, we conjecture that the upper bound can be improved to a double exponential bound.

The basic dependency pair method [1] forms the basis of our investigations. In particular we allow *argument filtering* for the dependency pairs. A similar result can be shown for *dependency graphs*, but we need to replace the triple exponential correspondence by an even faster growing (but still elementary) correspondence, as shown in the extended version of this paper [17]. Future work will concentrate on establishing better bounds for the studied variants, and analysing the derivational complexity induced by further refinements of the dependency pair method.

To summarise the contribution of this paper, we apply Theorem 4.2 to three well-studied simplification orders: LPO, MPO and KBO. Recall that the derivational complexity induced by LPO or MPO is multiple recursive or primitive recursive, respectively, cf. [20, 10]. Clearly these function classes are closed under elementary functions. Hence by Theorem 4.2 we obtain that the derivational complexity induced by the (basic) dependency pair method based on LPO (MPO) is multiple recursive (primitive recursive). On the other hand for a TRS $\mathcal{R}$ compatible with KBO we have that $\mathsf{dc}_{\mathcal{R}}$ belongs to $\mathsf{Ack}(\mathsf{O}(n), 0)$, cf. [14]. Thus applying the theorem in the context of KBO yields that the derivational complexity function induced by the dependency pair method based on KBO is majorised by the Ackermann function. Recall that in all three cases the bounds are tight (see [20, 10, 14]) and using the same examples, we obtain tightness of the here established bounds.

To conclude, we consider a version of the Ackermann function, introduced by Hofbauer [11] in a slightly simpler way, which we denote as $\mathcal{R}_6$.

$$\begin{array}{rlr} \mathsf{i}(x) \circ (y \circ z) \to \mathsf{f}(x, \mathsf{i}(x)) \circ (\mathsf{i}(\mathsf{i}(y)) \circ z) & & \mathsf{i}(x) \to x \\ \mathsf{i}(x) \circ (y \circ (z \circ w)) \to \mathsf{f}(x, \mathsf{i}(x)) \circ (z \circ (y \circ w)) & & \mathsf{f}(x, y) \to x \end{array}$$

Notice that $\mathcal{R}_6$ is not simply terminating and the derivational complexity of $\mathcal{R}_6$ dominates the Ackermann function. (The latter follows by the same argument as in [11].) However, termination can be shown easily by the (basic) dependency pair method in conjunction with argument filtering and KBO.

There are nine dependency pairs. For the argument filtering π, we set $\pi(\mathsf{f}) = \pi(\mathsf{f}^\sharp) = \pi(\mathsf{i}^\sharp) = 1$, $\pi(\mathsf{i}) = [1]$, and $\pi(\circ) = \pi(\circ^\sharp) = [1, 2]$. To apply Proposition 2.2 we use the reduction pair $(\geqslant^{\pi}_{\mathsf{KBO}}, >^{\pi}_{\mathsf{KBO}})$ induced by the (admissible) weight function w with $w_0 = 1$, $w(\circ) = w(\circ^\sharp) = 1$, and $w(\mathsf{i}) = 0$, together with the precedence $\mathsf{i} \succ \circ, \circ^\sharp$. Hence, by Theorem 4.2 the derivational complexity of $\mathcal{R}_6$ belongs to $\mathsf{Ack}(\mathsf{O}(n), 0)$ and this bound is optimal, compare [14].

Acknowledgements. The second author would like to thank Dieter Hofbauer for his hospitality, and fruitful discussions about a draft of this paper, during the his stay in Kassel.

References

[1] T. Arts and J. Giesl. Termination of term rewriting using dependency pairs. *TCS*, 236(1,2):133–178, 2000.

[2] F. Baader and T. Nipkow. *Term Rewriting and All That.* Cambridge University Press, 1998.

[3] Nachum Dershowitz. Termination dependencies. In *Proc. 6th WST*, Technical Report DSIC-II/15/03, Universidad Politecnica de Valencia, pages 27–30, 2003.

[4] J. Endrullis, J. Waldmann, and H. Zantema. Matrix interpretations for proving termination of term rewriting. *JAR*, 40(3):195–220, 2008.

[5] A. Geser. *Relative Termination*. PhD thesis, Universität Passau, 1990.

[6] N. Hirokawa and A. Middeldorp. Automating the dependency pair method. *IC*, 199 (1,2):172–199, 2005.

[7] N. Hirokawa and A. Middeldorp. Tyrolean termination tool: Techniques and features. *IC*, 205:474–511, 2007.

[8] N. Hirokawa and G. Moser. Automated complexity analysis based on the dependency pair method. In *Proc. 4th IJCAR*, volume 5195 of *LNCS*, pages 364–379, 2008.

[9] N. Hirokawa and G. Moser. Complexity, graphs, and the dependency pair method. In *Proc. 15th LPAR*, volume 5330 of *LNCS*, pages 652–666, 2008.

[10] D. Hofbauer. Termination proofs by multiset path orderings imply primitive recursive derivation lengths. *TCS*, 105(1):129–140, 1992.

[11] D. Hofbauer. *Termination Proofs and Derivation Lengths in Term Rewriting Systems*. PhD thesis, Technische Universität Berlin, 1992.

[12] D. Hofbauer and C. Lautemann. Termination proofs and the length of derivations. In *Proc. 3rd RTA*, volume 355 of *LNCS*, pages 167–177, 1989.

[13] A. Koprowski and J. Waldmann. Arctic termination ... below zero. In *Proc. 19th RTA*, volume 5117 of *LNCS*, pages 202–216, 2008.

[14] I. Lepper. Derivation lengths and order types of Knuth-Bendix orders. *TCS*, 269 (1,2):433–450, 2001.

[15] G. Moser. Derivational complexity of Knuth Bendix orders revisited. In *Proc. 13th LPAR*, volume 4246 of *LNCS*, pages 75–89, 2006.

[16] G. Moser, A. Schnabl, and J. Waldmann. Complexity analysis of term rewriting based on matrix and context dependent interpretations. In *Proc. 28th FSTTCS*, volume 08004 of *Dagstuhl Seminar Proceedings*, pages 304–315, 2008.

[17] Georg Moser and Andreas Schnabl. The derivational complexity induced by the dependency pair method. Technical report, University of Innsbruck, 2009. Available online at `http://cl-informatik.uibk.ac.at/users/aschnabl/`.

[18] TeReSe. *Term Rewriting Systems*, volume 55 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2003.

[19] R. Thiemann. *The DP Framework for Proving Termination of Term Rewriting*. PhD thesis, University of Aachen, 2007.

[20] A. Weiermann. Termination proofs for term rewriting systems with lexicographic path orderings imply multiply recursive derivation lengths. *TCS*, 139(1,2):355–362, 1995.

[21] H. Zantema. Termination of term rewriting by semantic labelling. *FI*, 24(1,2): 89–105, 1995.